

LIE COALGEBRAS AND RATIONAL HOMOTOPY THEORY II: HOPF INVARIANTS

DEV SINHA AND BENJAMIN WALTER

We give a new, definitive resolution of the homotopy periods problem modulo torsion. This problem was addressed most famously by Sullivan, on the first page and then Section 11 of his seminal paper [24]. There Sullivan starts to give explicit integrals over a sphere which determine whether two maps from that sphere are equivalent in the quotient of the homotopy groups of a simply connected space modulo their torsion subgroups. These integrals are a direct generalization of Whitehead’s formula for Hopf invariant [25], and they encode generalized linking numbers. More integrals along these lines were given by Haefliger [10] and some framework was developed by Novikov [17]. Before Sullivan’s work, Boardman and Steer [3] gave a set of linking number invariants defined for maps to a suspension. Finally, Hain in his thesis [11] uses Chen integrals as his model for the bar construction to give one resolution of the homotopy period question, but produces integrals over products of spheres with simplices rather than over spheres themselves and an auxiliary formalism is required to address redundancy.

We give a complete set of “integrals” (we can use integer-valued cochains if we wish) over the sphere which distinguish maps modulo torsion, proceeding from a simple use of the bar construction. We also connect these integrals to Quillen functors in the framework of differential graded Lie coalgebras, as developed in [23]. On the formal side, we are then able to evaluate Hopf invariants on Whitehead products and are able to understand the naturality of these maps in the long exact sequence of a fibration. For applications, we can show for example that the rational homotopy groups of homogeneous spaces are controlled by classical linking numbers.

We proceed in two steps, first using the classical bar complex to define integer-valued homotopy functionals which coincide with evaluation of the cohomology of ΩX on the looping of a map from S^n to X . One basic, apparently new, observation is that calculations in the classical bar complex yield the Hopf invariant formula of Whitehead (as well as those of Haefliger and Sullivan). We establish basic properties and give examples. In the second part we use the Harrison complex on commutative cochains, the standard cohomology theory for commutative algebras, and thus must switch to rational coefficients. Using our graph coalgebraic presentation, we show that a product-coproduct formula established geometrically in the bar complex descends to the duality predicted by Koszul-Moore theory.

In summary, we find that rational homotopy groups are controlled by “linking phenomena,” and that these linking phenomena are best described in the language of Lie coalgebras. One direction we plan to pursue further is the use of Hopf invariants to realize Koszul-Moore duality isomorphisms in general. A second direction we plan to pursue is that of spaces which are not simply connected, where the graph coalgebra models seem relevant even for $K(\pi, 1)$ spaces.

1. HOPF INVARIANTS FROM THE BAR COMPLEX.

In these first sections when dealing exclusively with the bar complex, we only need associative cochains and so work integrally. Later when dealing with the Lie coalgebraic model of a space, we switch to commutative cochains and work over the rationals. For consistency with historical practice we use $C^*(X)$ to denote the standard cochains with the cup product on a simplicial set X , or equivalently any subalgebra

1991 *Mathematics Subject Classification.* 55P62; 16E40, 55P48.

Key words and phrases. Hopf invariants, Lie coalgebras, rational homotopy theory, graph cohomology.

whose inclusion induces an isomorphism on cohomology. Similarly, we let $A^*(X)$ denote the PL forms on X or equivalently a subalgebra model. It is a funny accident of history that A^* is commutative while C^* is associative but not commutative (except in an E_∞ sense).

Definition 1.1. To set notation, let $B(R)$ denote the bar complex on an associative differential graded algebra R , defined in the standard way as the total complex of a bicomplex spanned by monomials $x_1 | \cdots | x_n$, where the x_i have positive degree, multilinear in each variable. The “internal” differential d_R is given by extending that of R by the Leibniz rule. We write d_μ for the “external” differential, defined by removing bars and multiplying. We write d_B for the total differential. If ω is a monomial in the bar complex, the internal degree of ω is the sum of the degrees of its component elements, and the weight of ω is its number of component elements. The total degree of ω is its internal degree minus its weight.

Let $B(X)$ denote $B(C^*(X))$, and let $H_B^*(X)$ denote $H_*(B(C^*(X)))$.

Throughout the paper, we generally suppress the suspension and desuspension operators s and s^{-1} , as where they need to appear is always determined by context. We may include them when for example they facilitate computing signs. Also, we will assume throughout that X is a simply-connected space.

The classical work of Adams-Hilton and Eilenberg-Moore established that $H_B^*(X)$ is isomorphic to the cohomology of the based loop space of X . We now show that the homology of the bar complex is also the natural setting for Hopf invariants. Topologically, we are passing from a map $f : S^n \rightarrow X$ to its looping $\Omega f : \Omega S^n \rightarrow \Omega X$, on which we evaluate cohomology classes from the bar complex. But the way in which we do the evaluation, and the properties we derive, have not to our knowledge been previously considered. We start with a standard calculation in the bar complex for the sphere. We include a proof because the central ingredient – weight reduction – generalizes to a method for explicit computation of Hopf invariants.

Lemma 1.2. $H_B^{n-1}(S^n)$ is rank one, generated by an element of weight one corresponding to the generator of $H^n(S^n)$.

Proof. Suppose $\alpha \in B^{n-1}(S^n)$ is a cycle. Since α has finitely many terms, its terms have finite maximal weight k . Write $\alpha|_k$ for the weight k terms of α . If $k > 1$ then its internal differential $d_{C^*}(\alpha|_k) = 0$, so $\alpha|_k$ gives a cocycle in $\otimes_k \bar{C}^*(S^n)$. By the Künneth theorem, $\otimes_k \bar{C}^*(S^n)$ has no homology in degree n . Thus $\alpha|_k$ is exact in $\otimes_k \bar{C}^*(S^n)$. Any choice of cobounding expression will determine a $\beta \in B(S^n)$ with $d_{C^*}\beta = \alpha|_k$. Therefore $\alpha - d_B\beta$ is a lower weight expression in $B^{n-1}(S^n)$ homologous to α . Inductively, we have that α is homologous to a cycle of weight one, so the map from $H^n(S^n)$ to $H_B^{n-1}(S^n)$ including the weight-one cocycles is surjective.

By applying the same weight-reduction argument to a cochain $\beta \in B^n(S^n)$ with $d\beta = x$ for x weight one, we see that the map from $H^n(S^n)$ to $H_B^{n-1}(S^n)$ is injective. $\square$

Definition 1.3. Let $\gamma \in B^{n-1}(S^n)$ be a cocycle. Define $\tau(\gamma) \simeq \gamma$ to be a choice of weight one cocycle to which γ is cohomologous.

Define $\int_{B(S^n)}$ to be the map from cocycles in $B^{n-1}(S^n)$ to $\mathbb{Z}$ given by $\int_{B(S^n)} \gamma = \int_{S^n} \tau(\gamma)$, where $\int_{S^n}$ denotes evaluation on the fundamental class of S^n .

From Lemma 1.2 it is immediate that the map $\int_{B(S^n)}$ is well defined and induces the isomorphism $H_B^{n-1}(S^n) \cong \mathbb{Z}$.

The standard way to use cohomology to define homotopy functionals is to pull back and evaluate. This is essentially all we do to define our generalized Hopf invariants, allowing through the definition of $\int_{B(S^n)}$ for a homology between the cocycle we pull back and one which we already know how to evaluate.

Definition 1.4. Define the Hopf pairing $\langle \cdot, \cdot \rangle_\eta : H_B^{n-1}(X) \times \pi_n(X) \rightarrow \mathbb{Z}$ by sending $[\gamma] \times [f]$ to $\int_{B(S^n)} f^*(\gamma)$.

We call $\tau(f^*(\gamma))$ the Hopf cochain (or form) of γ pulled back by f . We name the associated maps $\eta : H_B^{n-1}(X) \rightarrow \text{Hom}(\pi_*(X), \mathbb{Z})$ and $\eta^\dagger : \pi_*(X) \rightarrow \text{Hom}(H_B^{n-1}(X), \mathbb{Z})$. We say $\eta(\gamma) \in \text{Hom}(\pi_*(X), \mathbb{Z})$ is the Hopf invariant associated to γ .

The choice of Hopf cochain is not unique, but the corresponding Hopf invariants are. It is immediate that the Hopf invariants are functorial. Moreover, the definitions hold with any ring coefficients. Topologically we have the following interpretation.

Proposition 1.5. $\eta(\gamma)(f)$ coincides with the evaluation of the cohomology class given by γ in $H^{n-1}(\Omega X)$ on the image under Ωf of the fundamental class in $H_{n-1}(\Omega S^n)$.

1.1. Examples.

Example 1.6. A cocycle of weight one in $B(X)$ is just a closed cochain on X , which may be pulled back and immediately evaluated. Decomposable elements of weight one in $B(X)$ are null-homologous, consistent with the fact that products evaluate trivially on the Hurewicz homomorphism.

Example 1.7. Let ω be a generating 2-cocycle on S^2 and $f : S^3 \rightarrow S^2$. Then $\gamma = -\omega|_{\omega}$ is a cocycle in $B(S^2)$ which f pulls back to $-f^*\omega|f^*\omega$, a weight two cocycle of total degree two on S^3 . Because $f^*\omega$ is closed and of degree two on S^3 , it is exact. Let $d^{-1}f^*\omega$ be a choice of a cobounding cochain. Then

$$d(d^{-1}f^*\omega|f^*\omega) = f^*\omega|f^*\omega + (d^{-1}f^*\omega \smile f^*\omega).$$

Thus $f^*\gamma$ is homologous to $(d^{-1}f^*\omega \smile f^*\omega)$, and the corresponding Hopf invariant is $\int_{S^3} d^{-1}f^*\omega \smile f^*\omega$, which is the classical formula for Hopf invariant given by Whitehead [25], and generalized to maps from arbitrary domains by O'Neill [18]. To our knowledge, these formulae have not previously been related to the bar construction.

Expressions involving choices of d^{-1} for some cochains will be a feature of all of our formulae. As Sullivan points out when defining similar formulae in [24], on the sphere one can make this explicit, as in the proof of the Poincaré Lemma.

Example 1.8. Let $X = S^n \vee S^m$ and let x be a representative generator of $H^n(S^n)$ and similarly y on S^m . Then $\gamma = x|y$ is a cocycle in $B(X)$. Let $W : S^{n+m-1} \rightarrow S^n \vee S^m$ be the universal Whitehead product. More explicitly, let $p_1 : D^n \times D^m \rightarrow S^m$ be projection onto D^m followed by the canonical quotient map and let $p_2 : D^n \times D^m \rightarrow S^n$ be defined similarly. Decompose S^{n+m-1} as

$$S^{n+m-1} = \partial(D^n \times D^m) = S^{n-1} \times D^m \cup D^n \times S^{m-1}.$$

Then $W|_{S^{n-1} \times D^m} = p_1|_{S^{n-1} \times D^m}$ and $W|_{D^n \times S^{m-1}} = p_2|_{D^n \times S^{m-1}}$.

Proceeding as in the previous example, $\langle \gamma, W \rangle_\eta = (-1)^{|x|+1} \int_{S^{n+m-1}} d^{-1}W^*x \smile W^*y$. But note that these cochains extend to $D^n \times D^m$. Namely, $d^{-1}p_1^*x$ on $D^n \times D^m$ restricts to $d^{-1}W^*x$, and similarly W^*y is the restriction of p_2^*y . Now we evaluate as follows.

$$(-1)^{|x|+1} \int_{S^{n+m-1}} d^{-1}W^*x \smile W^*y = \int_{\partial(D^n \times D^m)} (d^{-1}p_1^*x \smile p_2^*y) = \int_{D^n \times D^m} p_1^*x \smile p_2^*y = \int_{D^n} x \cdot \int_{D^m} y = 1.$$

The change in sign in the first equality above is due to the change in orientation on the fundamental class induced by the isomorphism $S^{n+m-1} \cong \partial(D^n \times D^m)$, and the second equality is Stokes' Theorem.

We conclude that the Hopf invariant of γ detects the Whitehead product, which recovers a theorem from [10]. This first case of evaluation of a Hopf invariant on a Whitehead product will be generalized below.

Example 1.9. Let X be a manifold with a collection $\{W_i\}_{i=1}^k$ of proper oriented submanifolds of codimension two or greater, disjoint from each other and the basepoint. Denote the associated Thom classes by ω_i . Because the W_i are disjoint, $\gamma = \omega_1| \cdots | \omega_k$ is a cocycle in $B(X)$. If $f : S^d \rightarrow X$ where d is the total degree of γ (namely the sum of the codimensions of the W_i minus k) then one associated Hopf form is

$$\tau(f^*(\gamma)) = f^*\omega_1 \smile d^{-1}(f^*\omega_2 \smile d^{-1}(\cdots \smile (f^*\omega_{k-1} \smile d^{-1}f^*\omega_k) \cdots)).$$

Because f is based we can choose a diffeomorphism of $S^d \setminus \{\text{basepoint}\}$ with $\mathbb{R}^d$, and consider the submanifolds $f^{-1}W_i \subset \mathbb{R}^d$.

We construct $d^{-1}f^*\omega_k$ by taking the Thom class of the manifold given by extending $f^{-1}W_k$ upwards in $\mathbb{R}^d$ (say, in the first coordinate) a fixed large distance and “capping it off” once we have moved outside of a fixed neighborhood of the $f^{-1}W_i$ ’s. Then $f^*\omega_{k-1} \smile d^{-1}f^*\omega_k$ will be the Thom class of the submanifold of $f^{-1}W_{k-1}$ of points which lie above (in $\mathbb{R}^d$) some point of $f^{-1}W_k$. Proceeding in similar fashion, the Hopf invariant of f will be the generic count of collections of a point in $f^{-1}W_1 \subset \mathbb{R}^d$ which lies above a point in $f^{-1}W_2$ which in turn lies above a point in $f^{-1}W_3$, etc. This count is a generalized linking number, which specializes to a construction of Boardman and Steer [3]. It is equal to the degree of the Gauss map from $\prod_k f^{-1}W_i$ to $\prod_{k=1} S^{d-1}$ sending $(x_1, \dots, x_k)$ to $(\dots, \frac{x_i - x_{i-1}}{\|x_i - x_{i-1}\|}, \dots)$. For those familiar with the cohomology of ordered configuration spaces (see [21]), these values are encoding the image in homology of the fundamental class of $\prod f^{-1}(W_i)$ included in $\text{Conf}_k(\mathbb{R}^d)$.

Formulae for Hopf invariants may be interpreted in the language of intersection theory and linking behavior of submanifolds, which in some cases have boundary. Indeed, generalizing the definition of Hopf invariants through linking numbers was the starting point for our project, eventually leading to the formalism of Lie coalgebras.

Example 1.10. For an arbitrary X and cochains x_i, y_i and θ on X with $dx_i = dy_i = 0$ and $d\theta = \sum (-1)^{|x_i|} x_i \smile y_i$, the cochain $\gamma = \sum x_i |y_i + \theta \in B(X)$ is closed. The possible formulae for the Hopf invariant are all of the form

$$\langle \gamma, f \rangle_\eta = \int_{S^n} \left(f^*\theta - \sum \left((-1)^{|x_i|} t \cdot d^{-1}f^*x_i \smile f^*y_i + (1-t) \cdot f^*x_i \smile d^{-1}f^*y_i \right) \right),$$

for some real number t . This generalizes a formula given in the Computations section of [9] and is also present in [10].

By choosing $t = \frac{1}{2}$ we see that reversing the order to consider $\sum y_i |x_i$ will yield the same Hopf invariant, up to sign. Thus $\sum x_i |y_i \mp y_i |x_i$ yields a zero Hopf invariant. There are many Hopf invariants from the classical bar construction which are zero, a defect which will be remedied by using the Lie coalgebraic bar construction.

Example 1.11. In applications, Hopf forms are easily computed using a weight reduction technique introduced in the proof of Lemma 1.2. The bigrading of $B(X)$ is used as follows.

Suppose X is such that there is a weight three cocycle in $B(X)$ of the form

$$\gamma = x_1 |x_2 |x_3 - x_{12} |x_3 + x_{123},$$

where $dx_i = 0$, $dx_{12} = x_1 \smile x_2$, $dx_{123} = x_{12} \smile x_3$ and $x_2 \smile x_3 = 0$, x_1 has odd degree and x_2 has even degree. Consider

$$\alpha = d^{-1}f^*x_1 |f^*x_2 |f^*x_3 + d^{-1}(d^{-1}f^*x_1 \smile f^*x_2 - f^*x_{12}) |f^*x_3.$$

Here we observe that $(d^{-1}f^*x_1 \smile f^*x_2 - f^*x_{12})$ is closed and thus exact in order to know that we may find a d^{-1} in $C^*(S^n)$. We write down $d(\alpha) = (d_{C^*X} + d_\mu)(\alpha)$ in $B(S^n)$, which is naturally a second-quadrant bicomplex, as in Figure 1.

Observe that

$$f^*(\gamma) + d(\alpha) = (f^*x_{123} - d^{-1}(d^{-1}f^*x_1 \smile f^*x_2 - f^*x_{12}) \smile f^*x_3).$$

The right-hand side is a weight one cocycle in $B(S^n)$, meaning that it gives a Hopf cochain of γ pulled back by f .

$$\begin{array}{ccc}
& -f^*x_1|f^*x_2|f^*x_3 & \\
& \uparrow d_C & \\
d^{-1}f^*x_1|f^*x_2|f^*x_3 & \xrightarrow{d_\mu} & \begin{array}{l} (d^{-1}f^*x_1 \smile f^*x_2)|f^*x_3 \\ -(d^{-1}f^*x_1 \smile f^*x_2 - f^*x_{12})|f^*x_3 \end{array} \\
& \uparrow d_C & \\
d^{-1}(d^{-1}f^*x_1 \smile f^*x_2 - f^*x_{12})|f^*x_3 & \xrightarrow{d_\mu} & -d^{-1}(d^{-1}f^*x_1 \smile f^*x_2 - f^*x_{12}) \smile f^*x_3
\end{array}$$

FIGURE 1. Calculation of $d(\alpha)$ in $B(S^n)$.

1.2. The generalized Hopf invariant one question. From Proposition 1.5 and the Milnor-Moore theorem that the map from $\pi_*(\Omega X) \otimes \mathbb{Q} \rightarrow H_*(\Omega X; \mathbb{Q})$ is injective [15], we have the following.

Proposition 1.12. *If X is simply connected, the map $\eta_B : H_B^*(X; \mathbb{Q}) \rightarrow \text{Hom}(\pi_{*-1}(X), \mathbb{Q})$ is surjective. Thus, the map $\eta_B : H_B^*(X) \rightarrow \text{Hom}(\pi_{*-1}(X), \mathbb{Z})$ is of full rank.*

The question of the kernel of this map, which is large, is addressed in the next sections by using a different bar construction. The rational PL cochains on a simplicial set are commutative, so we should be taking a bar construction over the Koszul dual cooperad, namely the Lie cooperad, rather than associative cooperad. That is, the correct homology theory for a commutative algebra is Harrison/ André-Quillen homology.

The question of the cokernel of the integral map is a natural generalization of one of the most famous questions in the history of topology, namely the Hopf invariant one problem. The cokernel is trivial immediately for X an odd sphere and is $\mathbb{Z}/2$ for X an even sphere other than S^2 , S^4 and S^8 by Adams' celebrated result.

1.3. Evaluation on Whitehead products. Some properties of Hopf invariants, such as naturality, are immediate. A deeper result is to use the standard coproduct of $B(X)$ to evaluate Hopf invariants on Whitehead products.

Let $f : S^n \rightarrow X$ and $g : S^m \rightarrow X$, and recall the definition of the Whitehead product:

$$[f, g] : S^{n+m-1} \xrightarrow{W} S^n \vee S^m \xrightarrow{f \vee g} X.$$

We have $(f \vee g)^* = f^* + g^*$ on $C^*(X)$, so on $B(X)$ the map $(f \vee g)^*$ acts by $(f^* + g^*)$ on each component of a bar expression. For example,

$$\begin{aligned}
(f \vee g)^*(x_1 | x_2) &= (f^* + g^*)x_1 | (f^* + g^*)x_2 \\
&= f^*x_1 | f^*x_2 + f^*x_1 | g^*x_2 + g^*x_1 | f^*x_2 + g^*x_1 | g^*x_2
\end{aligned}$$

Below we will generally mean for f^* and g^* to be considered as maps $B(X) \rightarrow B(S^n \vee S^m)$ as in this example, omitting the inclusions of and projections onto wedge factors.

Theorem 1.13. *Let γ be a cocycle in $B^{n+m-2}(X)$ and $f : S^n \rightarrow X$, $g : S^m \rightarrow X$. Then*

$$\langle \gamma, [f, g] \rangle_\eta = \sum_j \langle \alpha_j, f \rangle_\eta \cdot \langle \beta_j, g \rangle_\eta \mp \langle \alpha_j, g \rangle_\eta \cdot \langle \beta_j, f \rangle_\eta$$

where $\Delta\gamma \simeq \sum_j \alpha_j \otimes \beta_j$, with all α_j and β_j closed, and $\mp$ is minus the Koszul sign induced by moving $s^{-1}\alpha_j$ past $s^{-1}\beta_j$.

Our convention is for $\langle \gamma, f \rangle_\eta = 0$ if $f : S^n \rightarrow X$ and $|\gamma| \neq n - 1$. The relation $\Delta\gamma \simeq \sum_j \alpha_j \otimes \beta_j$ with α_i and β_i closed follows from noting that $\Delta\gamma$ is closed in $B(X) \otimes B(X)$ and applying the Künneth isomorphism.

Proof. Note that $\langle \gamma, [f, g] \rangle_\eta = \langle (f \vee g)^*\gamma, W \rangle_\eta$. So we may compute the Hopf invariant in S^{n+m-1} by doing intermediate work in $S^n \vee S^m$. Recall $C^*(S^n \vee S^m) \cong C^*(S^n) \oplus C^*(S^m)$, which induces a component-bigrading on $B(S^n \vee S^m)$ as follows. If μ is a monomial in $B(S^n \vee S^m)$, we say that $w \in B(S^n)$ is an S^n component of μ if it is a maximal length subword with support only in S^n ; define S^m components of μ similarly. Bigrade $B(S^n \vee S^m)$ by the number of S^n components and the number of S^m components. If $a, b \in C^*(S^n \vee S^m)$ have disjoint supports then $a \smile b = 0$. Thus the differential of $B(S^n \vee S^m)$ preserves component-bigrading.

For example, let $a_i, b_i \in C^*(S^n \vee S^m)$ where the a_i have support in S^n and the b_i in S^m . Then $\mu = b_1|a_2|a_3|b_2|b_3|b_4$ has S^n component $a_2|a_3$ and S^m components b_1 and $b_2|b_3|b_4$. Thus μ has component bigrading $(1, 2)$.

Given a cocycle $\gamma \in B^{n+m-2}(X)$, the pullback $(f \vee g)^*\gamma \in B(S^n \vee S^m)$ splits as a sum of cocycles in each component-bigrading. By the Künneth theorem, $B(S^n \vee S^m)$ has no homology in degree $n + m - 2$ away from bigrading $(1, 1)$ and possibly $(1, 0)$ and $(0, 1)$. So all terms of $(f \vee g)^*\gamma$ not in these bigradings are exact. The terms in bigrading $(1, 0)$ and $(0, 1)$ are $f^*\gamma$ and $g^*\gamma$ which are exact after being pulled back to S^{n+m-1} by W .

Bigrading $(1, 1)$ itself splits as the sum of two subcomplexes – one where the S^n component comes first, and the other where the S^m component comes first. The subcomplex of bigrading $(1, 1)$ with the S^n component first is isomorphic to $B(S^n) \otimes B(S^m)$. The terms of $(f \vee g)^*\gamma$ in this subcomplex are

$$\sum_{\Delta\gamma = \sum_i a_i \otimes b_i} f^*(a_i)|g^*(b_i).$$

If $\sum_i a_i \otimes b_i \simeq \sum_j \alpha_j \otimes \beta_j$ then the cobounding expression in $B(X) \otimes B(X)$ will determine a homotopy between the above terms and $\sum_j f^*(\alpha_j)|g^*(\beta_j)$. By the Künneth theorem this is cohomologous to $\sum_j \tau(f^*\alpha_j)|\tau(g^*\beta_j)$. But by our work in example 1.8 the Hopf pairing of this is $\sum_j \langle \alpha_j, f \rangle_\eta \cdot \langle \beta_j, g \rangle_\eta$ as desired.

The terms of $(f \vee g)^*\gamma$ in bigrading $(1, 1)$ with the S^m component first are similar, but with a shift of sign at the final step due to the change in orientation induced by the homeomorphism $D^n \times D^m \cong D^m \times D^n$. $\square$

We will see that Hopf invariants completely determine the homotopy Lie algebra of X rationally. Through this theorem we can recover some of the integral Lie algebra structure as well.

This theorem implies for example that the Hopf invariant of a cocycle of the form $\alpha|\beta \mp \beta|\alpha$ evaluates trivially on all Whitehead products. Indeed, we saw in Equation 1.10 that this Hopf invariant is zero. Theorem 1.13 suggests a key for identifying bar expressions with vanishing Hopf invariants. If we define in the obvious way an “anti-commutative coproduct” on the bar construction, the submodule where this coproduct vanishes will by Theorem 1.13 have Hopf invariants which evaluate trivially on Whitehead products. Quotienting by this submodule yields the Lie coalgebraic cobar construction, which we develop next.

2. HOPF INVARIANTS FROM THE LIE COALGEBRA MODEL OF A SPACE, AND THEIR COMPLETENESS

We now switch to the rational, commutative setting and we replace the bar construction with Andre-Quillen homology. Instead of using the classical Harrison complex presentation of commutative Andre-Quillen homology, we use the larger graph complex $\mathcal{E}(A)$ introduced in [23]. As we will note in Proposition 2.4, through most of this section the reader may safely replace $\mathcal{E}(A)$ with the Harrison complex and graphs with bar expressions. The exception to this is Lemmas 2.16 and 2.18.

2.1. Lie coalgebraic bar construction.

Definition 2.1. Let V be a vector space. Define (ungraded) $\mathbb{E}(V)$ to be the quotient of $\mathbb{G}(V)$ by $\text{Arn}(V)$, where $\mathbb{G}(V)$ is the span of the set of oriented acyclic graphs with vertices labeled by elements of V modulo multilinearity in the vertices, and where $\text{Arn}(V)$ is the subspace generated by arrow-reversing and Arnold expressions:

$$\begin{array}{lcl}
 \text{(arrow-reversing)} & & \begin{array}{c} \text{graph 1} \end{array} + \begin{array}{c} \text{graph 2} \end{array} \\
 \text{(Arnold)} & & \begin{array}{c} \text{graph 3} \end{array} + \begin{array}{c} \text{graph 4} \end{array} + \begin{array}{c} \text{graph 5} \end{array}
 \end{array}$$

The diagrams represent the following operations:

- arrow-reversing:** Two graphs enclosed in dashed ellipses. The first graph has vertices a and b with an arrow from a to b . The second graph has vertices a and b with an arrow from b to a .
- Arnold:** Three graphs enclosed in dashed ellipses. The first graph has vertices a , b , and c with arrows $a \rightarrow b$ and $b \rightarrow c$. The second graph has vertices a , b , and c with arrows $a \rightarrow b$ and $c \rightarrow b$. The third graph has vertices a , b , and c with arrows $a \rightarrow c$ and $b \rightarrow c$.

Here a , b , and c are elements of V labeling vertices of a graph which could possibly have edges connectioning to other parts of the graph (indicated by the ends of edges abutting a , b , and c), which are not modified in these operations.

A Lie cobracket is defined as

$$]G[= \sum_{e \in G} (G_1^{\hat{e}} \otimes G_2^{\hat{e}} - G_2^{\hat{e}} \otimes G_1^{\hat{e}}),$$

where e ranges over the edges of G , and $G_1^{\hat{e}}$ and $G_2^{\hat{e}}$ are the connected components of the graph obtained by removing e , which points to $G_2^{\hat{e}}$. We actually require a graded version of $\mathbb{E}(V)$ given in Section 3 of [23].

Proposition 2.2. *If V is graded in positive degrees, then $\mathbb{E}(V)$ is isomorphic to the cofree Lie coalgebra on V .*

Indeed, if V and W are linearly dual, then the *configuration pairing* of [22, 23] can be used to define a perfect pairing between $\mathbb{E}(V)$ and the free Lie algebra on W . We use this graphical presentation of Lie coalgebras to give a new presentation of the Harrison complex.

Definition 2.3. Let A be a one-connected commutative differential graded algebra with differential d_A and multiplication μ_A . Define $\mathcal{G}(A)$ to be the total complex of the bicomplex $(\mathbb{G}(s^{-1}\bar{A}), d_A, d_\mu)$. Here $s^{-1}\bar{A}$ is the desuspension of the ideal of positive-degree elements of A , the “internal” differential d_A is given by cofreely extending that of A by the Leibniz rule, and the “external” differential $d_\mu(g) = \sum_e \mu_e(g)$ where up to sign $\mu_e(g)$ contracts the edge e in g , multiplying the elements of A labeling its endpoints to obtain the new label. (For the sign convention see Definition 4.5 of [23].) Let $\mathcal{E}(A)$, the Lie coalgebraic bar construction, be the quotient $\mathcal{G}(A)/\text{Arn}(s^{-1}\bar{A})$.

We define internal degree and weight of monomials in $\mathcal{E}(A)$ as the sum of the degrees of component elements and the number of vertices in the graph monomial. As before, total degree is internal degree minus the weight.

Define $\mathcal{E}(X)$ to be $\mathcal{E}(A^*(X))$ where $A^*(X)$ is, as in the introduction, a model for commutative cochains on X , and let $H_{\mathcal{E}}^*(X)$ denote $H^*(\mathcal{E}(X))$.

Recall that the Harrison chains $B_H(A)$ are constructed dually to Quillen's functor $\mathcal{L}$. The bar construction of a commutative algebra is a Hopf algebra under the shuffle product and the splitting coproduct. The Harrison complex is given by quotienting to Hopf algebra indecomposables. Proposition 3.21 of [23] leads to the following.

Proposition 2.4. *There is a short-exact sequence of bicomplexes, giving an isomorphism of final terms*

$$\begin{array}{ccccccc} 0 & \longrightarrow & \text{Sh}(A) & \longrightarrow & B(A) & \longrightarrow & B_H(A) \longrightarrow 0 \\ & & \downarrow & & \phi \downarrow & & \cong \downarrow \\ 0 & \longrightarrow & \text{Arn}(A) & \longrightarrow & \mathcal{G}(A) & \longrightarrow & \mathcal{E}(A) \longrightarrow 0, \end{array}$$

where ϕ sends the bar expression $a_1|a_2|\cdots|a_n$ to the graph $\begin{array}{c} \nearrow^{a_2} \searrow^{a_n} \\ a_1 \end{array}$. Furthermore, this vertical isomorphism is an isomorphism of Lie coalgebras when the Harrison complex is given the Lie coalgebra structure defined by Schlessinger and Stasheff [20].

The presentation of Harrison homology via graphs has a critical advantage over the classical construction. While the set of generators of $\mathcal{G}(A)$ is larger than that of $B(A)$, the set of relations $\text{Arn}(A)$ are simpler to write down and are defined locally, unlike the shuffle relations $\text{Sh}(A)$, which is convenient for calculations and can be critical for some arguments. By abuse we will sometimes use bar notation to refer to elements of $\mathcal{E}(A)$, suppressing ϕ .

The significance of Harrison homology in this setting is that it bridges the worlds of cohomology and homotopy. One of the main thrusts of this paper is to give a new geometric understanding of this bridge through Hopf invariants. By a celebrated theorem of Barr [2], the quotient map $p : BA \rightarrow B_H A$ has a splitting e when working over a field of characteristic 0. Harrison homology is a special case of André-Quillen homology [1, 19] for one-connected commutative differential graded algebras. For convenience, we use Barr's splitting and the development of Hopf invariants for the standard bar complex to quickly establish basic properties. Deeper properties will require the Lie coalgebraic formulation.

2.2. Hopf invariants. As we did for the bar complex, we start with the fact that $H_{\mathcal{E}}^{n-1}(S^n)$ is rank one, generated by an element of weight one.

Definition 2.5. Given a cocycle $\gamma \in \mathcal{E}^{n-1}S^n$ we let $\tau(\gamma) \simeq \gamma$ be any cohomologous cocycle of weight one.

Write $\int_{\mathcal{E}(S^n)}$ for the map from cocycles in $\mathcal{E}^{n-1}(S^n)$ to $\mathbb{Q}$ given by $\int_{\mathcal{E}(S^n)} \gamma = \int_{S^n} \tau(\gamma)$.

Lemma 2.6. *The map $\int_{\mathcal{E}(S^n)}$ is well defined and induces the isomorphism $H_{\mathcal{E}}^n(S^n) \cong \mathbb{Q}$.*

Proof. We observe that $\int_{\mathcal{E}(S^n)} = \int_{B(S^n)} \circ e$, since Barr's splitting e is an isomorphism which is an "identity" on the weight one parts of these complexes. Thus this lemma follows from Lemma 1.2. $\square$

Remark 2.7. In light of our use of the classical bar construction in this step, one might view that construction as more fundamental. What seems to be the most fundamental bar construction is the graph coalgebraic bar construction $\mathcal{G}(A)$. We choose to use the classical bar construction only because it is more familiar and directly tied to cohomology of loopspaces. Logically, it would be better to stay within the context of graph coalgebras for explicit calculation and Lie coalgebras for formal properties, omitting reference to the standard bar construction and thus associative coalgebras altogether.

We can now take our Definition 1.4 of Hopf invariants and replace the bar complex everywhere by $\mathcal{E}$ (and the integers by the rational numbers). When necessary, we distinguish the Hopf pairings and Hopf invariants by decorations such as $\langle , \rangle_{\eta}^{\mathcal{E}}$ vs. $\langle , \rangle_{\eta}^B$, or $\eta^{\mathcal{E}}$ vs. η^B , etc.

Theorem 2.8. *Hopf invariants are compatible with the quotient map p and Barr's splitting e between $B(X)$ and $\mathcal{E}(X)$. That is, $\eta^B = \eta^{\mathcal{E}} \circ p$ and $\eta^{\mathcal{E}} = \eta^B \circ e$.*

Proof. We consider the adjoint maps and fix an $f \in \pi_n(X)$. The images of $(\eta^B)^\dagger(f)$ and $(\eta^\mathcal{E})^\dagger(f)$ are then given by the composites going from left to right in the following diagram.

$$\begin{array}{ccccc}
 B(X) & \xrightarrow{f^*} & B(S^n) & \xrightarrow{f_{B(S^n)}} & \mathbb{Q} \\
 \downarrow p & \uparrow e & \downarrow p & \uparrow e & \\
 \mathcal{E}(X) & \xrightarrow{f^*} & \mathcal{E}(S^n) & \xrightarrow{f_{\mathcal{E}(S^n)}} & \mathbb{Q}
 \end{array}$$

The result follows immediately from commutativity of the two squares (one involving p and one involving e) and two triangles in the above diagram. The squares commute due to naturality of the quotient map and Barr's splitting, and the triangles commute by our comment in the proof of Lemma 2.6. $\square$

We now see that all of the shuffles in the kernel of the quotient map $p : B(X) \rightarrow \mathcal{E}(X)$ give rise to the zero homotopy functional, constituting a large kernel.

Using this theorem, we can translate some theorems from the bar setting, such as the following corollary of Theorem 1.13.

Corollary 2.9. *Let γ be a cocycle in $\mathcal{E}^{n+m-2}(X)$ and $f : S^n \rightarrow X$, $g : S^m \rightarrow X$. Then*

$$\langle \gamma, [f, g] \rangle_\eta^\mathcal{E} = \sum_j \langle \alpha_j, f \rangle_\eta^\mathcal{E} \cdot \langle \beta_j, g \rangle_\eta^\mathcal{E}$$

where $[\gamma] \simeq \sum_j \alpha_j \otimes \beta_j$, with all α_j and β_j closed.

By recasting our Hopf invariants in terms of Lie coalgebras, we get some immediate applications.

Example 2.10. Consider a wedge of spheres $X = \bigvee_i S^{d_i}$. Let ω_i represent a generator of $H^{d_i}(S^{d_i})$ and W be the differential graded vector space spanned by the ω_i with trivial products and differential. Then W is a model for cochains on X and $\mathcal{E}(W) = \mathbb{E}(W)$ is just the cofree Lie coalgebra on W .

Theorem 1.13 along with Proposition 3.6 of [23] imply that the value of some Hopf invariant η_γ on an iterated Whitehead product P of the basic inclusions $\iota_i = [S^{d_i} \hookrightarrow X] \in \pi_{d_i}(X)$ is given by the configuration pairing (defined in Corollary 3.5 of [23]) of γ and P . This immediately implies that the ι_i generate a copy of a free Lie algebra in the rational homotopy groups of X and gives a complete understanding of what the Hopf invariants of any element of that free Lie algebra are.

Because this example is universal, a corollary to Theorem 1.13 is that in general the value of a Hopf invariant on an iterated Whitehead product can be calculated through a configuration pairing.

2.3. Completeness. Once we focus on $\mathcal{E}(X)$, the resulting Hopf invariants give a complete and sharp picture of rational homotopy groups.

Theorem 2.11. $\eta^\mathcal{E} : H_\mathcal{E}^*(X) \rightarrow \text{Hom}(\pi_*(X), \mathbb{Q})$ is an isomorphism.

There are three possible lines of proof. Using Theorem 1.13 and Example 2.10 we could show that η agrees with the isomorphism given by the Quillen equivalences established in [23]. Alternately we could try to start with the surjectivity of this map, which follows from Proposition 1.12, and try to calculate the kernel. Instead we shall prove more directly that η is an isomorphism by developing a long exact sequence of a fibration for $\mathcal{E}$. This approach yields an independent proof and gives a more direct geometric understanding of the isomorphism in Theorem 2.11.

Definition 2.12. Let $\mathcal{E}(E, F)$ be the kernel of the map from $\mathcal{E}(E)$ to $\mathcal{E}(F)$ induced by inclusion. Define relative Hopf invariants for $\pi_n(E, F)$, by pulling back a cycle in $\mathcal{E}(E, F)$ to $\mathcal{E}(D^n, S^{n-1})$ and then reducing to weight one and evaluating.

That $H_{\mathcal{E}}^n(D^n, S^{n-1})$ is rank one generated by a cycle of weight one follows from a direct calculation. To see that relative Hopf invariants are well defined we may use the same weight-reduction argument.

By construction, there is a long exact sequence for $H_{\mathcal{E}}^*(F)$, $H_{\mathcal{E}}^*(E)$ and $H_{\mathcal{E}}^*(E, F)$. Hopf invariants give a pairing between between this long exact sequence and the long exact sequence for relative homotopy groups. For clarity, we will drop the identifier η from our Hopf pairings below and instead subscript them by their ambient space.

Proposition 2.13. *The following long exact sequences pair compatibly*

$$\begin{aligned} \cdots \longrightarrow \pi_n(F) \xrightarrow{i_*} \pi_n(E) \xrightarrow{j_*} \pi_n(E, F) \xrightarrow{\partial} \pi_{n-1}(F) \longrightarrow \cdots \\ \cdots \longleftarrow H_{\mathcal{E}}^n(F) \xleftarrow{i^*} H_{\mathcal{E}}^n(E) \xleftarrow{j^*} H_{\mathcal{E}}^n(E, F) \xleftarrow{\delta} H_{\mathcal{E}}^{n-1}(F) \longleftarrow \cdots \end{aligned}$$

That is, for $f \in \pi_n(F)$ and $\gamma \in H_{\mathcal{E}}^n(E)$, $\langle i^*\gamma, f \rangle_F = \langle \gamma, i_*f \rangle_E$, and similarly elsewhere.

Proof. For i_* and i^* this is immediate from naturality, as both $\langle i^*\gamma, f \rangle_F$ and $\langle \gamma, i_*f \rangle_E$ are equal to $\int_{S^n} \tau((f \circ i)^*\gamma)$. The argument for $\langle j^*\gamma, f \rangle_E$ and $\langle \gamma, j_*f \rangle_{(E, F)}$ is the same once one substitutes $\pi_n(E, pt.)$ for $\pi_n(E)$ and $H_{\mathcal{E}}^n(E, pt.)$ for $H_{\mathcal{E}}^n(E)$. In this case one now sees both pairings as given by $\int_{D^n} \tau((f \circ j)^*\gamma)$.

Finally we consider the connecting homomorphisms. By definition $\delta : H_{\mathcal{E}}^{n-1}(F) \rightarrow H_{\mathcal{E}}^n(E, F)$ proceeds by taking a cocycle γ , lifting it at the cochain level to $\mathcal{E}^{n-1}(E)$ and then taking its coboundary. So $\langle \delta(\gamma), f \rangle_{(E, F)} = \int_{D^n} \tau(f^*d_{\mathcal{E}}\bar{\gamma})$, where $\bar{\gamma}$ is an extension of γ to all of E . We have that $\langle \gamma, \partial f \rangle_F = \int_{\partial D^n} \tau(f^*(\gamma))$. Restricted to ∂D^n , $\tau(f^*d_{\mathcal{E}}\bar{\gamma})$ can be chosen to be $d\tau(f^*(\gamma))$, so Stokes' theorem yields the result. $\square$

We now connect with the exact sequences of a fibration. Let $p : E \rightarrow B$ be a fibration and write $\lambda : \pi_n(B) \rightarrow \pi_n(E, F)$ for the standard map defined by lifting.

Proposition 2.14. *The map $p^* : \mathcal{E}(B) \rightarrow \mathcal{E}(E, F)$ is a weak equivalence.*

Corollary 2.15. *Let γ be a cycle in $\mathcal{E}(E, F)$. Then γ is homologous to $p^*(\bar{\gamma})$ for some $\bar{\gamma}$ in $\mathcal{E}B$.*

Proof of Proposition 2.14. We use the maps induced by the Sullivan model for the fiber inclusion map i , as given in [8] §15(a):

$$\begin{array}{ccccc} A_{PL}F & \xleftarrow{i^*} & A_{PL}E & \xleftarrow{p^*} & A_{PL}B \\ \uparrow m_F \simeq & & \uparrow m \simeq & \nearrow \iota & \\ (\Lambda V_F, \bar{d}) & \xleftarrow{\varepsilon} & (A_{PL}B \otimes \Lambda V_F, d) & & \end{array}$$

Here $(\Lambda V_F, \bar{d})$ is the Sullivan model for $A_{PL}F$, ι is the map induced by the inclusion of the unit $\mathbb{Q} \rightarrow \Lambda V_F$, and ε is the map induced by the augmentation $A_{PL}B \rightarrow \mathbb{Q}$.

Write $\hat{\mathcal{E}}(E, F)$ for the kernel of the map $\varepsilon : \mathcal{E}(A_{PL}B \otimes \Lambda V_F, d) \rightarrow \mathcal{E}(\Lambda V_F, \bar{d})$. There is an induced quasi-isomorphism $\hat{\mathcal{E}}(E, F) \xrightarrow{\simeq} \mathcal{E}(A_{PL}E, A_{PL}F)$. Our work is completed in the lemmas which follow by establishing that the inclusion maps i_1 and i_2 in the diagram below are both quasi-isomorphisms, and therefore so is the induced map on kernels $\mathcal{E}A_{PL}B \rightarrow \hat{\mathcal{E}}(E, F)$.

$$\begin{array}{ccccc} \mathcal{E}A_{PL}F & \xleftarrow{i^*} & \mathcal{E}A_{PL}E & \xleftarrow{\delta} & \mathcal{E}(E, F) \\ \uparrow \varepsilon m_F \simeq & & \uparrow \varepsilon m \simeq & \nearrow p^* & \uparrow \simeq \\ \mathcal{E}(\Lambda V_F) & \xleftarrow{\varepsilon_*} & \mathcal{E}(A_{PL}B \otimes \Lambda V_F) & \xleftarrow{\bar{\delta}} & \hat{\mathcal{E}}(E, F) \\ \uparrow i_1 \simeq & & \uparrow i_2 \simeq & \nearrow \iota_* & \uparrow \simeq \\ V_F & \xleftarrow{pr} & \mathcal{E}A_{PL}B \oplus V_F & \xleftarrow{\delta} & \mathcal{E}A_{PL}B \end{array}$$

Here the differentials on V_F and ΛV_F are the standard restrictions of that on $A_{PL}B \otimes \Lambda V_F$.

□

An alert reader might note that while we have generally steered clear of the traditional approach to rational homotopy through minimal models, we do use the Sullivan model for a fibration here. However, we really only needed that the cochains of a total space of a fibration is a twisted tensor product of those on the fiber and base, and Sullivan model theory is the most convenient and well-known reference for this. We could just as well have used earlier work of Brown [5] giving this result for simplicial cochains of fibrations, along with a translation to PL cocahins and the easy fact that we can make free models (we did not require minimality).

Lemma 2.16. *The inclusion $V_F \rightarrow \mathcal{E}(\Lambda V_F)$ sending v to $\begin{smallmatrix} v \\ \bullet \end{smallmatrix}$ is a quasi-isomorphism.*

Proof. The proof proceeds in a similar manner as that of Theorem 19.1 of [8]. The homogeneous elements of $\mathcal{E}\Lambda V_F$ are graphs whose vertices are decorated by nonempty words in V_F , the sum of whose lengths we call the total word length. We define a map of complexes $h : \mathcal{E}\Lambda V_F \rightarrow \mathcal{E}\Lambda V_F$ which away from the image of V_F gives a chain homotopy between the identity map and a map which increases the weight of a graph, while keeping its total word length fixed if possible or which is zero if that is not possible. Since weight is bounded by total word length, iterated applications of this will eventually produce a null-homology of any cycle not in V_F .

Write a generic long graph $\gamma \in \mathcal{E}^{m+1}\Lambda V$ of total word length n as

$$\gamma = \left(\begin{array}{c} v_{k_1} \otimes \cdots \quad v_{k_j} \otimes \cdots \otimes v_{k_{j+l}-1} \quad v_{k_m} \otimes \cdots \otimes v_n \\ \nearrow \quad \searrow \quad \nearrow \quad \searrow \quad \nearrow \\ v_1 \otimes \cdots \quad \cdots \quad \cdots \quad \cdots \quad \cdots \end{array} \right).$$

Define $h(\gamma)$ to be zero if the total word length of γ is equal to $m+1$ (in which case $\gamma = v_1 | \cdots | v_m$). For γ with at least one word of length two decorating some vertex, the expression $h(\gamma)$ is a sum over the letters in such words of new graphs obtained by removing each letter in turn and using it to decorate a new vertex attached to the vertex containing the previous word while fixing the rest of the graph. Explicitly,

$$h(\gamma) = \frac{1}{n} \sum_{\substack{j \text{ with} \\ k_{j+1} - k_j > 1}} \sum_{k_j \leq p \leq k_{j+1} - 1} (-1)^{\kappa(p)} \left(\begin{array}{c} v_p \\ \downarrow \\ v_{k_j} \otimes \cdots \otimes \widehat{v_p} \otimes \cdots \otimes v_{k_{j+1}-1} \\ \nearrow \quad \searrow \\ \cdots \quad \cdots \end{array} \right)$$

where the $(-1)^{\kappa(p)}$ is the Koszul sign coming from moving a degree one operator to v_p 's vertex and moving v_p to the front of its word and across an s^{-1} .

More generally if γ is any decorated graph, we define h as a sum over the letters in the words of length at least two decorating the vertices of γ , removing each letter in turn and using it to decorate a new vertex as above. The internal ordering of the vertices in the resultant graph is the same as for γ with the split off vertex occuring immediately before the vertex it was taken from.

That h is well defined follows easily from the local nature of the anti-symmetry and Arnold relations in the definition of $\mathcal{E}$.

Away from V_F , h decreases degree by one (due to the unwritten s^{-1} in front of v_p) and increases graph length by one. It is straightforward to check that $d_{\mathcal{E}}h + hd_{\mathcal{E}} = id + (\text{graphs of greater length})$ outside of V_F . Note that the Arnold and arrow reversing relations are required in order to get this equality for graphs with internal vertices decorated by singleton words. In particular on the basic case of graphs whose total word length is $m+1$, h is a chain homotopy between the identity map and the projection onto V_F . □

Remark 2.17. It is not clear how to define h on the bar complex presentation of the Harrison complex, and it seems that doing so would be difficult because the shuffle relations in the Harrison complex are not local.

The proof presented above is cleaner than the standard proof of the dual result in Proposition 22.8 of [8], which requires going through the universal enveloping algebra and then invoking Theorem 19.1 of [8].

Lemma 2.18. *The inclusion $\mathcal{E}A_{PL}B \oplus V_F \rightarrow \mathcal{E}(A_{PL}B \otimes \Lambda V_F)$ is a quasi-isomorphism.*

Sketch of proof. The proof is similar to the previous lemma. Homogeneous elements of $\mathcal{E}(A_{PL}B \otimes \Lambda V_F)$ are graphs whose vertices are decorated by an element of $A_{PL}B$ and a word in V_F . On any vertex, the decorating element from $A_{PL}B$ may be trivial or the V_F word may be empty, but not both at once.

As in the proof of Lemma 2.16, we use a chain homotopy h which evaluates on a graph as a sum over replacements of each vertex “pulling out” V_F letters in turn and connecting them back to the vertex by a new edge. As before it is simple to check that $d_{\mathcal{E}}h + hd_{\mathcal{E}} = id + (\text{graphs of greater length})$ outside of $\mathcal{E}A_{PL}B \oplus V_F$ $\square$

Remark 2.19. To our knowledge, the dual of this result does not appear in the literature. Furthermore, attempting to prove the dual statement by methods analogous to the proof of Proposition 22.8 in [8] would be difficult.

Proposition 2.20. *Evaluation of Hopf invariants is compatible with the isomorphisms $\pi_n(E, F) \cong \pi_n(B)$ and $H_{\mathcal{E}}^n(E, F) \cong H_{\mathcal{E}}^n(B)$. That is, $\langle p^*\gamma, f \rangle_{(E, F)} = \langle \gamma, p_*f \rangle_B$ and $\langle \gamma, \lambda f \rangle_{(E, F)} = \langle \bar{\gamma}, f \rangle_B$.*

Proof. The first equality is immediate since both $\langle p^*\gamma, f \rangle_{(E, F)}$ and $\langle \gamma, p_*f \rangle_B$ are equal to $\int_{D^n} \tau((p \circ f)^*\gamma)$.

For the second equality, we combine Corollary 2.15 with some elementary identifications as follows,

$$\langle \gamma, \lambda f \rangle_{(E, F)} = \int_{D^n} \tau((\lambda f)^*\gamma) = \int_{D^n} \tau((\lambda f)^*p^*\bar{\gamma}) = \int_{D^n} \tau((p \circ \lambda f)^*\bar{\gamma}) = \int_{D^n} \tau(f^*\bar{\gamma}) = \langle \bar{\gamma}, f \rangle_B.$$

$\square$

We can now quickly prove the main theorem.

Proof of Theorem 2.11. Applying the perfect pairing between the $\mathcal{E}$ and homotopy long exact sequences of a fibration to the Postnikov tower of a space, it is enough to know that $\eta^{\mathcal{E}}$ gives an isomorphism on Eilenberg-MacLane spaces.

Recall that $A_{PL}K(\mathbb{Q}^m, n)$ is quasi-isomorphic to a free graded algebra with m generators in degree n . Lemma 2.16 gives a null homotopy of all elements of $\mathcal{E}(\Lambda_s^n \mathbb{Q}^m)$ except for these generators. The Hopf invariants associated to these generators, given by simply pulling back cohomology and evaluating, are linearly dual to $\pi_n(K(\mathbb{Q}^m, n))$ by the Hurewicz theorem. $\square$

This proof required the existence but not the uniqueness of the Postnikov tower. On the whole, the only theorems from topology which this theory of Hopf invariants requires are this existence and the fact that a fibration can be modeled by a twisted tensor product of cochain models. The rest of the basic input has been algebraic, namely in the development of the Lie cooperadic bar construction.

3. APPLICATIONS AND OPEN QUESTIONS

We give applications and present some questions. We have listed one such already, namely the generalized Hopf invariant one question. The results of the previous section gives rise to a new approach to both the classical question and its generalization. Namely one could try to understand how the classical bar construction over the integers, modulo Harrison shuffles, fails to yield an exact sequence of a fibration.

3.1. Wedges of spheres. Revisiting Example 2.10, recall that a model for cochains on a wedge of spheres X is just the graded vector space W with one generator w_i in the appropriate dimension for each sphere, with trivial products and differential. Theorem 2.11 now implies that the linear dual of $\pi_*(X)$ is the cofree Lie coalgebra on W . Using Corollary 2.9 and the perfect pairing between free Lie algebras and cofree Lie coalgebras to deduce that the homotopy groups of X are just the free Lie algebra on W , yielding a short and conceptually simple route to the rational Hilton-Milnor Theorem. Moreover, we not only know what the homotopy Lie algebra is but see explicitly how to determine whether two maps into wedges of spheres are homotopic. The vector space $\mathcal{E}(W)$ is spanned by cocycles of the form $w_{\sigma(1)}|w_{\sigma(2)}|\cdots|w_{\sigma(k)}$ as σ varies over the symmetric group. Following Example 1.9 represent w_i by the “Thom class” associated to a point p_i (away from the basepoint) of each wedge factor S^{d_i} . Given some map $f : S^n \rightarrow X$ we let $W_i = f^{-1}(p_i) \subset \mathbb{R}^{d_i}$. The argument in Example 1.9 applies, and we interpret the Hopf invariant of γ_σ as a linking number of the W_i , counting the number of times (with signs) that a point in $W_{\sigma(1)}$ lies “directly above” one in $W_{\sigma(2)}$ which in turn lies above one in $W_{\sigma(3)}$, etc. Thus the rational homotopy class of f is determined by such counts.

3.2. Homogeneous spaces. If G is a Lie group, then it is rationally homotopy equivalent to a generalized Eilenberg-MacLane space. It is formal, and its rational cohomology algebra is graded free on odd-dimensional generators. By Lemma 2.16, $\mathcal{E}(G) \simeq V$, where V is the vector space of generators of the cohomology ring included as weight one cocycles. The associated Hopf invariants, which are complete if G is simply-connected, are just evaluation of cohomology classes, reflecting the standard fact that the homotopy groups of an Eilenberg-MacLane space map isomorphically to their Hurewicz image.

Now consider the homogeneous space G/H and the standard Puppe sequence (for fibrations, as in [16]) $H \xrightarrow{i} G \rightarrow G/H \xrightarrow{f} BH \xrightarrow{Bi} BG$, which gives rise to an exact sequence when applying $H_\mathcal{E}^*$. We apply Lemma 2.16 for H , G , BH and BG to see that $H_\mathcal{E}^*$ of each of these is a vector space (which we denote $V_{BH} \cong sV_H$, $V_{BG} \cong sV_G$) of generators of their cohomology rings, or more precisely of indecomposables. Thus, $H_\mathcal{E}^*(G/H)$ is isomorphic to the direct sum of the kernel and suspended cokernel of $i^* : V_G \rightarrow V_H$. Hopf invariants associated to the cokernel of i^* are simple to interpret. A cohomology class in V_{BH} can be pulled back to G/H and evaluated on homotopy, and it will be trivial if were in turn pulled back from BG . For the kernel, if $x \in V_G$ and $i^*(x) = 0$ then $i^*(sx) = 0$ in V_{BH} . By definition this means that $Bi^*(x)$ is decomposable and thus equal to $\sum a_i b_i$ for some collection of $a_i, b_i \in H^*(BH)$. Then $\sum f^*(a_i)|f^*(b_i)$ will be a cocycle in $\mathcal{E}(G/H)$, whose Hopf invariant we associate to x .

Thus, the Hopf invariants for G/H , which are complete if G/H is simply connected, are given by either evaluation of cohomology pulled back from BH or by classical linking invariants associated to cocycles of BH whose products are trivial when pulled back to G/H . Thus while weight one cocycles account for the Hopf invariants of a generalized Eilenberg-MacLane space, weight two cocycles were required for these spaces which have a two-stage Postnikov system. In general the highest weight which appears in $H_\mathcal{E}^*(X)$ should reflect the number of stages the smallest tower atop which X sits whose fibers are generalized Eilenberg-MacLane spaces.

3.3. Configuration spaces. Next consider the ordered configuration space $\text{Conf}_n(\mathbb{R}^d)$, with $n = 3$. Its cohomology algebra is generated by classes a_{12}, a_{13} and a_{23} where for example a_{12} is pulled back from the map which sends (x_1, x_2, x_3) to $\frac{x_2 - x_1}{\|x_2 - x_1\|} \in S^{d-1}$. These cohomology classes satisfy the Arnold identity $a_{12}a_{23} + a_{23}a_{31} + a_{31}a_{12} = 0$ (the same identity at the heart of the graphical approach to Lie coalgebras, an overlap which we explain below). Its homotopy Lie algebra is generated by classes whose images in homology are dual, which we denote b_{ij} , for which “ x_i and x_j orbit each other.” Their Lie brackets satisfy the identity that $[b_{12}, b_{23}] = [b_{23}, b_{31}] = [b_{31}, b_{12}]$.

The best way to construct Hopf invariants is not with the standard a_{ij} classes since their products are non-trivial. The simplest cocycle in $\mathcal{E}(\text{Conf}_3(\mathbb{R}^d))$ using these classes would be

$$\begin{array}{c} a_{23} \\ \nearrow \\ a_{12} \end{array} + \begin{array}{c} a_{31} \\ \nearrow \\ a_{23} \end{array} + \begin{array}{c} a_{12} \\ \nearrow \\ a_{31} \end{array} + \bullet, \quad \theta$$

where θ is a cochain cobounding the Arnold identity. Consider instead the submanifold of points (x_1, x_2, x_3) which are collinear. This submanifold has three components, which we label by which x_i is “in the middle.” These are proper submanifolds, so we consider their Thom classes, which we denote Co_i . By intersecting these with the cycles representing the homology generators b_{ij} we see that for example Co_1 is cohomologous to $a_{31} + a_{12}$. Because they are disjoint, we have that any graph $\begin{array}{c} Co_j \\ \nearrow \\ Co_i \end{array}$ is a cocycle in $\mathcal{E}(\text{Conf}_n(\mathbb{R}^d))$. We may use Corollary 2.9 to see for example that

$$\begin{aligned} \left\langle \begin{array}{c} Co_2 \\ \nearrow \\ Co_1 \end{array}, [b_{12}, b_{23}] \right\rangle_\eta &= (a_{31} + a_{12})(b_{12}) \cdot (a_{12} + a_{23})(b_{23}) + (a_{12} + a_{23})(b_{12}) \cdot (a_{31} + a_{12})(b_{23}) \\ &= 1 \cdot 1 + 1 \cdot 0 = 1. \end{aligned}$$

Thus, to understand an element f of $\pi_{2d-3}(\text{Conf}_3(\mathbb{R}^d))$ it suffices to understand the linking behavior of $f^{-1}(Co_i)$. This calculation is reflected in the results of [6], which used relative Hopf invariants of a “Gauss map” for knots to give a new interpretation of the simplest finite-type knot invariant. Having a framework to detect iterated Whitehead products in this context is a central motivation for development of these Hopf invariants.

3.4. The graph complex. Though we relied on the classical bar complex, from the point of view of proofs using $\mathcal{E}(X)$, the graph complex is a much more natural home for the development of Hopf invariants. Basic constructions and proofs proceed in the same manner as those for the bar complex, with some arguments applying verbatim. The product-coproduct formula also goes through, using graph-cutting and maximal connected components in place of the standard coproduct and components in the bar complex. Since the quotient map $\mathcal{G}(S^n) \rightarrow \mathcal{E}(S^n)$ is the identity on weight one, the proof of Theorem 2.8 applies to show that $\eta^{\mathcal{G}} = \eta^{\mathcal{E}} \circ p$. However, if we wished to use $\eta^{\mathcal{G}}$ to show that $\eta^{\mathcal{E}}$ is well defined we would either need an alternate, direct, proof that Hopf invariants vanish on arrow reversing and Arnold expressions, or else we would need a splitting of the quotient map $\mathcal{G}(A) \rightarrow \mathcal{E}(A)$. In modern terminology, the splitting of the quotient $B(A) \rightarrow \mathcal{E}(A)$ is given by the first Eulerian idempotent $e_*^{(1)}$ ([13] §4.5), which together with the other Eulerian idempotents, this splits the bar complex into the Hodge decomposition. It is tempting to seek a similar decomposition of the graph complex. Such a decomposition might give an alternate way of understanding the Eulerian idempotents.

4. COMPARISON WITH OTHER APPROACHES

As mentioned in the introduction, our theorems give a resolution of the homotopy periods problem, of explicitly representing homotopy functionals. There have been a variety of approaches, and the present theory of Hopf invariants connects well with them. A formula for homotopy periods using the same ingredients, namely pulling back forms to the sphere, taking d^{-1} and wedge products, and then integrating, was featured in Sullivan’s seminal paper [24]. We have not been able to compare our formula with Sullivan’s, since that formula comes from a minimal model for path spaces which uses a chain homotopy which is also a derivation. In some examples, we cannot find such a chain homotopy, and if one relaxes the condition of being a derivation then the inductive formula is not clear.

In the same vein, some of the lower weight cases of our constructions were treated in work of Haefliger [10] and Novikov [17]. Haefliger also gives formulae for evaluating these forms on Whitehead products, which now follows from Theorem 1.13. Haefliger’s comment was that “It is clear that one could continue this way for higher Whitehead products, if one is not afraid of complicated formulas.” The bracket-cobracket formalism and the configuration pairing make these formulae simple, at least conceptually. Haefliger uses

Hopf invariants associated to cocycles in $\mathcal{E}(X)$ of the form $x_1 | \cdots | x_n$ where $x_i x_{i-1} = 0$ to show that such X have summands of the corresponding free Lie algebra in their homotopy groups, which is also immediate from the present approach.

Hain's thesis [11] addresses the rational homotopy period using Chen integrals. Here one appeals to the Milnor-Moore theorem that identifies rational homotopy within loop space homology [15], so that dually loop space cohomology must give all homotopy functionals. The main work is to find an irredundant collection, and Hain's formalism to do so does not immediately connect to Quillen functors and Whitehead product structure. The geometric heart of Chen integrals are the evaluation maps $\Delta^n \times \Omega X \rightarrow X^n$ which evaluates a loop at n points. Thus when writing down these functionals explicitly, the integrals are over $S^n \times \Delta^k$ not S^n itself.

As mentioned in our Example 1.9, our Hopf invariants also connect with those of Boardman and Steer [3, 4]. Their Hopf invariants are maps $\lambda_n : [\Sigma Y, \Sigma X] \rightarrow [\Sigma^n Y, \Sigma X^{\wedge n}]$. In some cases, the latter group can be computed, yielding homotopy functionals. A key intermediate they take is defining a map $\mu_n : [\Sigma Y, \bigvee_{i=1}^n X_i] \rightarrow [\Sigma^n Y, \bigwedge_{i=1}^n X_i]$. Comparing the proof of Theorem 6.8 of [4] with our Example 1.9 we see that their functionals agree with ours in this setting. While their technique is more general in the direction of analyzing maps out of any suspension as opposed to only spheres, our approach allows analysis of homotopy groups of spaces which are not suspensions. Indeed, the behavior of $\mathcal{E}(\Sigma X)$ is fairly trivial since the bar complex collapses.

Finally, consider Example 2.10 which showed that the value of a Hopf invariant on a Whitehead product is given by the configuration pairing between free Lie algebras and coalgebras. This pairing appears in the homology and cohomology of ordered configuration spaces. Recall from Example 1.9 that if $\{W_i\}$ is a disjoint collection of proper submanifolds of X and f is a map $S^d \rightarrow X$, then Hopf invariants are encoding the map on homology induced by inclusion of $\prod f^{-1}(W_i)$ in $\text{Conf}_k(\mathbb{R}^d)$. As Fred Cohen often says, "this cannot be a coincidence." In this case the connection is readily explained through Cohen's work on the homology of iterated loop spaces.

Consider $Y = \bigvee_{i=1}^k S^{d_i+2}$ and let $n = d_1 + \cdots + d_k + k - 1$, the dimension in which one can see a Whitehead product P of the inclusion maps ι_i with each map appearing once. Then $\pi_n(Y) \cong \pi_{n-2}(\Omega^2 \Sigma^2 (\bigvee S^{d_i}))$ which maps to $H_{n-2}(\Omega^2 \Sigma^2 (\bigvee S^{d_i}))$ under the Hurewicz homomorphism. Loop space theory identifies this homology with that of $\bigcup_m \text{Conf}_m(\mathbb{R}^2) \wedge_{\Sigma_m} (S^{d_1} \vee \cdots \vee S^{d_k})^{\wedge m}$, which in turn has a summand coming from the homology of $\text{Conf}_k(\mathbb{R}^2) \times_{\Sigma_k} (\bigcup_{\sigma \in \Sigma_k} S^{d_{\sigma(1)}} \wedge \cdots \wedge S^{d_{\sigma(k)}})$. Since the Σ_k action is free on the union of products of spheres, we get that this is simply

$$H_{n-2}(\text{Conf}_k(\mathbb{R}^2) \wedge S^{\sum d_i}) \cong H_{k-1}(\text{Conf}_k(\mathbb{R}^2)) \cong \mathcal{L}ie(n).$$

(See [21] for an expository treatment of this last isomorphism.) In Remark 1.2 of [7], established at the end of Section 13, Cohen gives a diagram establishing the compatibility of Whitehead products and Browder brackets, through the Hurewicz homomorphism. This result implies that a Whitehead product P maps to this $\mathcal{L}ie(k)$ summand in the canonical way, going to the homology class with the same name (up to sign).

In Example 2.10, we showed that the Hopf invariant η_γ evaluated on P according to the configuration pairing between free Lie algebras and coalgebras, modeled in the operad/cooperad pair $\mathcal{L}ie$ and $\mathcal{E}il$. Thus a cohomology class $g \in \mathcal{E}il(k) \cong H^{k-1}(\text{Conf}_k(\mathbb{R}^2))$ represented by a graph (see [21]) gets mapped under the linear dual of the Hurewicz homomorphism to essentially the same graph (with each vertex label replaced by a volume form on the corresponding sphere) in $\mathcal{E}(Y)$.

In summary, for wedges of spheres (and other double-suspensions) our Hopf invariants coincide with applying the Hurewicz homomorphism and then evaluating on the cohomology of configuration spaces, which explains the combinatorial similarity between our Hopf invariants and that cohomology. At one point we outlined a proof of Corollary 2.9 using this kind of loop space machinery, but we later found the more elementary approach given here.

REFERENCES

1. Michel André, *Homologie des algèbres commutatives*, Springer-Verlag, Berlin, 1974, Die Grundlehren der mathematischen Wissenschaften, Band 206. MR MR0352220 (50 #4707)
2. Michael Barr, *Harrison homology, Hochschild homology and triples*, J. Algebra **8** (1968), 314–323. MR MR0220799 (36 #3851)
3. J. M. Boardman and B. Steer, *Axioms for Hopf invariants*, Bull. Amer. Math. Soc. **72** (1966), 992–994. MR MR0202138 (34 #2012)
4. ———, *On Hopf invariants*, Comment. Math. Helv. **42** (1967), 180–221. MR MR0221503 (36 #4555)
5. Edgar H. Brown, Jr., *Twisted tensor products. I*, Ann. of Math. (2) **69** (1959), 223–246. MR MR0105687 (21 #4423)
6. Ryan Budney, James Conant, Kevin P. Scannell, and Dev Sinha, *New perspectives on self-linking*, Adv. Math. **191** (2005), no. 1, 78–113. MR MR2102844 (2005h:57013)
7. Frederick R. Cohen, Thomas J. Lada, and J. Peter May, *The homology of iterated loop spaces*, Springer-Verlag, Berlin, 1976. MR MR0436146 (55 #9096)
8. Yves Félix, Stephen Halperin, and Jean-Claude Thomas, *Rational homotopy theory*, Graduate Texts in Mathematics, vol. 205, Springer-Verlag, New York, 2001. MR MR1802847 (2002d:55014)
9. Phillip A. Griffiths and John W. Morgan, *Rational homotopy theory and differential forms*, Progress in Mathematics, vol. 16, Birkhäuser Boston, Mass., 1981. MR MR641551 (82m:55014)
10. André Haefliger, *Whitehead products and differential forms*, Differential topology, foliations and Gelfand-Fuks cohomology (Proc. Sympos., Pontificia Univ. Católica, Rio de Janeiro, 1976), Lecture Notes in Math., vol. 652, Springer, Berlin, 1978, pp. 13–24. MR MR505648 (80a:55012)
11. Richard M. Hain, *Iterated integrals and homotopy periods*, Mem. Amer. Math. Soc. **47** (1984), no. 291, iv+98. MR MR727818 (85d:55015)
12. D. K. Harrison, *Commutative algebras and cohomology*, Trans. Amer. Math. Soc. **104** (1962), 191–204. MR MR0142607 (26 #176)
13. Jean-Louis Loday, *Cyclic homology*, second ed., Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], vol. 301, Springer-Verlag, Berlin, 1998, Appendix E by María O. Ronco, Chapter 13 by the author in collaboration with Teimuraz Pirashvili. MR MR1600246 (98h:16014)
14. Walter Michaelis, *Lie coalgebras*, Adv. in Math. **38** (1980), no. 1, 1–54. MR MR594993 (82b:17016)
15. John W. Milnor and John C. Moore, *On the structure of Hopf algebras*, Ann. of Math. (2) **81** (1965), 211–264. MR MR0174052 (30 #4259)
16. Yasutoshi Nomura, *On mapping sequences.*, Nagoya Math J. **17** (1960), 111–145. MR MR521692 (80h:55016)
17. S. P. Novikov, *Analytical theory of homotopy groups*, Topology and geometry—Rohlin Seminar, Lecture Notes in Math., vol. 1346, Springer, Berlin, 1988, pp. 99–112. MR MR970074 (89k:55027)
18. Edward J. O’Neill, *Higher order Massey products and links*, Trans. Amer. Math. Soc. **248** (1979), no. 1, 37–66. MR MR521692 (80h:55016)
19. Daniel Quillen, *On the (co-) homology of commutative rings*, Applications of Categorical Algebra (Proc. Sympos. Pure Math., Vol. XVII, New York, 1968), Amer. Math. Soc., Providence, R.I., 1970, pp. 65–87. MR MR0257068 (41 #1722)
20. Michael Schlessinger and James Stasheff, *The Lie algebra structure of tangent cohomology and deformation theory*, J. Pure Appl. Algebra **38** (1985), no. 2-3, 313–322. MR MR814187 (87e:13019)
21. Dev P. Sinha, *Operads and knot spaces*, J. Amer. Math. Soc. **19** (2006), no. 2, 461–486 (electronic). MR MR2188133
22. ———, *A pairing between graphs and trees.*, math.QA/0502547, 2006.
23. Dev P. Sinha and Ben Walter, *Lie coalgebras and rational homotopy theory, I*, math.AT/0610437.
24. Dennis Sullivan, *Infinitesimal computations in topology*, Inst. Hautes Études Sci. Publ. Math. **47** (1977), 269–331 (1978). MR MR0646078 (58 #31119)
25. J. H. C. Whitehead, *An expression of Hopf’s invariant as an integral*, Proc. Nat. Acad. Sci. U. S. A. **33** (1947), 117–123. MR MR0020255 (8,525h)

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OREGON, EUGENE, OR 97403

E-mail address: dps@math.uoregon.edu

MIDDLE EAST TECHNICAL UNIVERSITY, NORTHERN CYPRUS CAMPUS, GÜZELYURT, MERSIN 10, TURKEY.

E-mail address: benjamin@metu.edu.tr